*Journal of Immunological Methods*, 166 (1993) 303–304
Elsevier Science Publishers B.V.



# Contents to volume 166 (1993)

**Number 1 – 5 November 1993**

*Research reports*

Detection and quantification of secreted soluble FcγRIIA in human sera by an enzyme-linked immunosorbent assay
*A. Astier, H. De la Salle, J. Moncuit, M. Freund, J.-P. Cazenave, W.-H. Fridman, D. Hanau and J.-L. Teillaud (France)* 1

Analysis of lymphocyte aggregation using digital image analysis
*L.L. Munn, M.W. Glacken, B.W. McIntyre and K. Zygourakis (USA)* 11

Human IgG preparations isolated by ion-exchange or protein G affinity chromatography differ in their glycosylation profiles
*A. Bond, M.G. Jones and F.C. Hay (UK)* 27

Rapid quantification of C3a and C5a using a combination of chromatographic and immunoassay procedures
*H. Hartmann, B. Lübbers, M. Casaretto, W. Bautsch, A. Klos and J. Köhl (Germany)* 35

Rapid flow cytometric assay for the assessment of natural killer cell activity
*L. Chang, G.A. Gusewitch, D.B.W. Chritton, J.C. Folz, L.K. Lebeck and S.L. Nehlsen-Cannarella (USA)* 45

Covalent coupling of a steroid to microwell plates for use in a competitive enzyme-linked immunosorbent assay
*S. Yonezawa, A. Kambegawa and S. Tokudome (Japan)* 55

Immortalization of mouse intestinal epithelial cells by the SV40-large T gene. Phenotypic and immune characterization of the MODE-K cell line
*K. Vidal, I. Grosjean, J.-P. Revillard, C. Gespach and D. Kaiserlian (France)* 63

Analysis of active antibody concentration. Separation of affinity and concentration parameters.
*R. Karlsson, L. Fägerstam, H. Nilshans and B. Persson (Sweden)* 75

The concentrations of glutamine and ammonia in commercially available cell culture media
*S. Heeneman, N.E.P. Deutz and W.A. Buurman (Netherlands)* 85

A sensitive colorimetric assay for the release of tryptase from human lung mast cells in vitro
*S.E. Lavens, D. Proud and J.A. Warner (UK, USA)* 93

A capillary electrophoresis-based assay for the binding of $Ca^{2+}$ and phosphorylcholine to human C-reactive protein
*N.H.H. Heegaard and F.A. Robey (USA)* 103

Carrier determination for X-linked agammaglobulinemia using X inactivation analysis of purified B cells
*L.A. Alterman, M. De Alwis, S. Genet, R. Lovering, H. Middleton-Price, G. Morgan, A. Jones, S. Malcolm, R.J. Levinsky and C. Kinnon (UK)* 111

Purification and properties of immunotoxins containing one vs. two deglycosylated ricin A chains
*V. Ghetie, E. Swindell, J.W. Uhr and E.S. Vitetta (USA)* 117

Rapid immunofiltration assay of Newcastle disease virus using a silicon sensor
*W.E. Lee, H.G. Thompson, J.G. Hall, R.E. Fulton and J.P. Wong (Canada)* 123

Measurement of kinetic binding constants of viral antibodies using a new biosensor technology
*J.L. Pellequer and M.H.V. Van Regenmortel (France)* 133

$Ca^{2+}$ response in single human T cells induced by stimulation of CD4 or CD8 and interference with CD3 stimulation
*Y. Katayama, S. Miyazaki, Y. Oshimi and K. Oshimi (Japan)* 145

*Notes to authors* 155

**Number 2 – 3 December 1993**

*Research reports*

Rapid in vitro micro-cytotoxicity tests for the detection and quantitation of neutralizing antibodies to both viruses and toxins
*C. Van de Water, E.A. Van Dura, J.G.M.M. Van der Stap, R. Brands and W.J.A. Boersma (Netherlands)* 157

In vitro immunization of mouse spleen cells for the production of monoclonal IgG1 antibodies using an antigen-specific T helper cell clone (D.10.G4.1)
*S. Uthoff and T. Böldicke (Germany)* 165

A simple method for the evaluation of receptor binding capacity of modified cytokines
*M. Chianelli, A. Signore, R. Hicks, R. Testi, M. Negri and P.C.L. Beverley (Italy, UK)* 177

Measurement of eosinophil cationic protein (ECP) and eosinophil protein X/eosinophil derived neurotoxin (EPX/EDN). Time and temperature dependent spontaneous release in vitro demands standardized sample processing
*C.M. Reimert, L.K. Poulsen, C. Bindslev-Jensen, A. Kharazmi and K. Bendtzen (Denmark)* 183
The assessment of different methods of data reduction for qualitative assays
*J. Bishop, F.D.J. Dunstan, B.J. Nix and A. Swift (UK)* 191
Similar frequencies and kinetics of cytokine producing cells in murine peripheral blood and spleen. Cytokine detection by immunoassay and intracellular immunostaining
*B. Sander, I. Höidén, U. Andersson, E. Möller and J.S. Abrams (Sweden, USA)* 201
Measurement of membrane bound IgD and IgM on B lymphocytes
*M. Zhang, T. Brunnée, J. Kleine-Tebbe, K. Graf, A. O'Connor and G. Kunkel (Germany)* 215
An improved model for the examination of biological effects of locally administered cytokines
*F.M. Rolhwagen, N.D. Pacheco and S. Baqar (USA)* 223
Evidence of surface antigen detachment during incubation of cells with immunomagnetic beads
*C.P. Rubbi, D. Patel and D. Rickwood (UK)* 233
A fluorescence-based quantitative adhesion assay to study interactions between *Entamoeba histolytica* and human enterocytes. Effect of proinflammatory cytokines
*L. Flores-Romo, K.B. Bacon, T. Estrada-Garcia, M. Shibayama, V. Tsutsumi and A. Martinez-Palomo (Switzerland, Mexico)* 243
Quantitative analysis of platelet-activating factor in human plasma. Application to patients with liver cirrhosis and disseminated intravascular coagulation
*J. Sugatani, M. Miwa, Y. Komiyama and T. Murakami (Japan)* 251
New procedure for the detection of complement deficiency by ELISA. Analysis of activation pathways and circumvention of rheumatoid factor influence
*G.N. Fredrikson, L. Truedsson and A.G. Sjöholm (Sweden)* 263
Liposomes as vaccine carriers. Incorporation of soluble and particulate antigens in giant vesicles
*S.G. Antimisiaris, P. Jayasekera and G. Gregoriadis (UK)* 271

*Short communications*
The role of pH in modified ELISA procedures used for the estimation of functional antibody affinity
*D. Goldblatt, L. Van Etten, F.J. Van Milligen, R.C. Aalberse and M.W. Turner (UK, Netherlands)* 281
A vector driving the expression of foreign cDNAs in the MHC class II-positive cells of transgenic mice
*V. Kouskoff, H.-J. Fehling, M. Lemeur, C. Benoist and D. Mathis (France)* 287

*Letter to the editors*
Selective destruction of contaminating cells in monolayer cultures
*E.W. Johnstone and K.A. Williams (Australia)* 293

*Software notices*
TITRECAL: an MS-DOS program for automated calculation of antibody titres from ELISA data
*S.A. Tremain (Australia)* 295
ELEASY: a program for processing experimental enzymoimmunoassay data
*A. Herráez (Spain)* 297
Software archive. Current contents (August 25, 1993)
*S.J. Merrill (USA)* 299

*Corrigendum*
Technical influences on immunophenotyping by flow cytometry. The effect of time and temperature of storage on the viability of lymphocyte subsets. (J. Immunol. Methods 164 (1993) 263)
*T. Ekong, E. Kupek, A. Hill, C. Clark, A. Davies and A. Pinching (UK)* 301

*Contents to volume 166 (1993)* 303

*Notes to authors* 305